

African Research Review

An International *Multidisciplinary Journal, Ethiopia*

Vol. 2 (3), August, 2008

ISSN 1994-9057 (Print)

ISSN 2070--0083 (Online)

Identification of Allelochemicals from Terminalia Chebula (Pp. 306 -314)

Manikandan, M. & Rejula M.

Abstract

This study was an experiential research to identify the chemicals contained in terminalia chebula. The dried leaves and barks were ground to a fine powder in a Wiley Mill (40 mesh), using this powder for aqueous extract prepared by the method of Heisey (1990). Bioassay studies were also carried out following the method of Heisey (1990). The extraction of phenolic compounds for GC analysis was carried out by the method of Kil and Yum (1983). The comparison of the leaf and bark extracts was done. The result of our study showed that the leaf extracts of the T.chebula contains more quantity of hydroquinone, trans-cinnamic acid, gentisic acid, vanillic acid, syringic acid and transferulic acid (phenolic acids) otherwise known as allelochemicals than the bark and were more inhibitory to seedling growth of Cassia occidentals and Crotalaria retusa than the bark extracts in bioassay studies. The implication is that Terminalia Chebula will be extremely useful in future to control the weed growth in agroecosystem.

Key Words: *Terminnlia chebula, Cassia occidentals, Crotalaria retusa* and allelo chemicals.

Introduction

Plants produce a large variety of secondary products containing hydroxylated aromatics rings these substance are classified as allelochemicals, most of which are synthesized from phenylalanine, a

product of the shikimic acid path way (Taiz and Zeiger 1991). Phenolic acids in aerial parts, roots of allopathic crops actively exude phenolics such as caffeic, chlorogenic, isochlorogenic, p-coumaric, p-hydroxy benzoic and ferulic acids were detected in alfalfa root exudates and vegetative residues (Abdul Rahman and Habib 1989). These secondary metabolic products may be released in to environment, generally the rhizosphere in sufficient quantities to affect neighboring plants. Mostly research on allelopathy has been focused on the effect on interactions among weed species (Willison and Rice, 1968, Rasmussen and Rice 1971). The objectives of this study were to:

- (i) Develop a method of weed control by using allelopathic plant
- (ii) Quantify the allelopathic potential of *Terminalia Chebula* on weeds
- (iii) Identify the plant parts that are the most important source of allelopathic substances
- (iv) Identification of phenolic acids (Allelochemicals)

Methodology

Mature fresh leaves and barks of the *Terminalia chebula* were collected from Srivilliputtur Reserve Forest, Tamil Nadu, India. Leaves and barks were dried in an oven at $60^{\circ}\text{c} \pm 2^{\circ}\text{c}$ for four days powdered (40 mesh) and used for phenolic acid extraction and bioassay studies. Seeds of the *Cassica occidentalis* and *Crotalaria retusa* (weeds) were collected from Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu, India. The dried leaves and barks were ground to a fine powder in a Wiley Mill (40 mesh), using this powder for aqueous extract prepared by the method of Heisey (1990).

Preparation of Aqueous Extract

Ten grams of leaf and bark powder dissolved separately in 100 ml of distilled water in a beaker and kept for 24 hr at room temperature ($28^{\circ}\text{C} \pm 2^{\circ}\text{C}$) with occasional swirling. The solution was filtered through what man N0 1 filter paper and the volume was made up to 100 ml with distilled water. The aqueous extract was diluted with water to get 5, 10, 15 and 20% concentrations. The dilutions corresponded to 0.5, 0.1, 0.15 and 0.2% of water extractable materials.

Bioassay Studies

Bioassay studies were carried out following the method of Heisey (1990). Ten weed seeds were placed on what man N0 1 filter paper in petriplates (9cm x 2cm). Petriplates were moistened with 2ml/plate of leaf /bark aqueous extract/distilled water (control), and incubated in dark $28^{\circ}\text{C} \pm 2^{\circ}\text{C}$. The experimental design was a randomized complete block with five replicated for each treatment and control. Germination percentage, plumule and radicle lengthly were measured after three days.

Extraction of Phenolic Acid for GC Analysis

The extraction of phenolic compounds for GC analysis was carried out by the method of Kil and Yum (1983). 2 grams of powdered leaf and bark samples were defatted using hexane. Then the defatted powder was mixed with 100 ml of 70% methanol: 70% acetic acid (20%) and centrifuged at 3,000 rpm for 5 minutes. The resulting supernatant was saved and concentrated to 20 ml. The P^{H} of the supernatant was adjusted to 2.0 with 6N HCL. Then this solution was centrifuged at 3,000 rpm for 5min the residues were discarded and the supernatant was mixed and partitioned with 20 ml of hexane: distilled water (1:1) 5 times. The aqueous solution was saved and it was extracted with equal volume of ethanol: ethyl acetate (1:1) for 5 times. The resulting ethanol: ethyl acetate solution was concentrated and used for the analysis of phenolic acid by GC.

Phenolic acids were identified GC (HewlettPackard-5890USA) using J& W fused silica capillary and SE-54 column (30m x 0.2mm) column temperature was at 90°C (5min) 4°C/min 250 °C (20min). Injector and detector temperature was 280°C. The head pressure of column was 30Psi and split ratio was 36:1 the detector was FID the volume of the sample injected was 0.5-1.0 µl. Comparing the retention times of the peaks with those of commercial Sigma and Aldrich phenolic acid samples were used for the identification of each peak.

Results and Discussion

In bioassay studies of *T. chebula* on *C. occidentalis* the germination of plumule and radical length was decreased in all concentrations. In the highest concentration this inhibition was maximum (Table 1 and 2). The inhibitory effect on the seed germination by leaf aqueous extracts was gradual while the bark extract of *T. chebula* drastically inhibited the seed germination. A maximum of 28% reduction in germination was observed at 20% leaf extract

The reduction in plumule length by aqueous leaf extracts was gradual but the bark extracts showed rapid decreases in plumule length. The maximum of 22% and 60% reduction in plumule length was recorded at 20% leaf and bark extracts respectively.

A similar pattern of decrease was observed in radicle length of *C. occidentalis* seeds treated with leaf and bark aqueous extracts maximum of 22% and 42% reduction in radical length was noticed at 20% concentration of leaf and bark respectively

The aqueous leaf and bark extracts of *T. chebula* were inhibitory to the seed germination, plumule and radical length of *C. retusa* seeds. The maximum inhibitory effect was observed at the highest concentration (Table 3 and 4). The reduction in seed germination was gradual by the treatment of leaf aqueous extract of *T. chebula*, whereas the reduction was rapid by the bark aqueous extracts treatment. The maximum

reduction of 28% was recorded at 20% leaf and bark aqueous extract concentration respectively.

The inhibition of plumule length by the aqueous bark extract was more than the leaf aqueous extracts. At the highest concentration both the leaf and bark aqueous extracts of *T. chebula* showed 22% and 60% reduction in plumule length respectively. The reduction in radicle length was 22% at 20% leaf aqueous extract, while at the same concentration there was only 42% reduction observed in radicle length in bark aqueous extract.

Identification of phenolic acid

The GC analysis of phenolic acids in bark and leaf of *Terminalia chebula* showed the presence of following phenolic acids:

- Phenolic acid in *T.chebula* leaf: Hydroquinone (27.7ug/grams), trans-cinnamic acid (20.3ug/grams), genetic acid (153.8 ug/g), vanillic acid (34.4 u/grams), syringic acid (11.7ug/grams) and trans-ferulic acid (67.2ug/gram)
- Phenolic acid in *T.chebula* bark: Hydroquinone (15.8ug/g), salicylic acid (40.3 ug/g), genetic acid (3.2ug/g), vanillic acid (4.2ug/g), proto catechuic acid (5.7ug/g), syringic acid (49.2ug/g), p-coumaric acid (30.1 ug/g) trans ferulic acid (4.2 ug/g) and caffeic acid (2.1 ug/g).

The result of our study showed that the leaf extracts of the *T.chebula* were more inhibitory to seed germination in the plants tested than the bark extracts in bioassay studies. Heisey (1990) has reported similar results. He observed that the leaflets among the various plant parts of *Ailanthus altissima* showed the highest inhibitory effect on seed germination several weed and crop species. Leaf extracts of several trees like *Grewia oppositifolia*, *Ficus roxburghi* and *Bauhinia variegata* tested by kaletha et.al (1996) showed a higher rate of inhibition on the germination of maize, cowpea, finger millet and

soybean. Since the quantities of allelochemicals vary among different plant tissues (Putnam and Duke, 1978 and rice, 1984) the higher rate of inhibition by leaf extracts may be explained by the higher amount of allelochemicals present in them. In fact the content of Phenolic compounds was higher in leaf extracts than the bark extracts as is evident from *T.chebula* Phenolic acid studies.

Conclusion

P-coumaric acid has been reported (Lee and Monsi, 1963) as the important allelochemicals from red pine and it has been implicated in respiration and Ca^{++} ion uptake and consequently in inhibition of mung bean hypocotyls length (Demos et al. 1975) syringic acid showed a significant reduction in mung bean hypocotyls length with out inhibiting the respiration and Ca^{++} uptake and it inhibited the root elongation in wheat, rye and mung bean (Vaughan and Ord, 1991) above this evidence the inhibition of radical, plumule and seed germination may due to the presence of these phenolic acids. These allelochemicals also found in the leaf of *T.chebula* is inhibitory to seedling growth of *Cassia occidentalis* and *Crotalaria retusa* than the bark extracts in bioassay studies. The implication is that *Terminalia Chebula* will be extremely useful in future to control the weed growth in agroecosystem.

References:

- Abdul Rahman, A.A. and Habib, S.A. (1989). Allelopathic effect of alfalfa (*Medicago sativa*) on blady grass (*Imperat cylindrica*) *Journal of Chemical Ecology*.15: 2289-2300.
- Al-saadawi, I.S. and Rice, E.L. (1982). Allelopathic effect of *polygonum aviculare* L. II. Isolation, characterization and biological activities of phytotoxins. *Journal of Chemical Ecology*.8: 1011-23
- Demos, E.L., Woolwin, M, Wilson, R.H. and Mcmillan, C. (1975). The effect of ten phenolic compounds on hypocotyls growth and mitochondria metabolism of mung bean. *American Journal of Botany*.62: 97-102.

- Heisey, R.M. (1990). Allelopathic and herbicidal effects of extracts from tree of heaven (*Alianthus altissima*) *American Journal of Botany*.77: 662-670.
- Kaletha, M.S., Bhatt, B.P and Todatria, N.P. (1996). Tree-crop interaction in traditions in traditional agro forestry systems of Grahewal Himalaya.I. Phytotoxic effects of farm trees on food crops. *Allelopathy Journal* 3:247-250.
- Kil, B.S.and Y.J., Yim. (1983): Allelopathic effects of *Pinus densiflora* on under growth of red pine forest. *Journal of Chemical Ecology*.9:1135-1151.
- Lee, I.K. and Monsi, M. (1963). Ecological studies on *Pinus densiflora* forest .I. effect of plant substances on the floristic composition of the under growth. *BotanyMagazine*.76: 400-413.
- Putnam, A.R. and Duke, W.B. (1978). Allelopathy in agro ecosystems. *Annual.Review of Phyto pathology*. 16:431-451.
- Rasmussen, J.A and Rice, E.L. (1971). Allelopathic effect of *Sporobolus pyramidatus* on vegetation patterning. *American Midland Naturalist*, 86:309-326.
- Tang, C.S and young, C.C. (1982). Collection and identification of allelopathic compounds from the undisturbed root system of *Bigalta limporgtass* (*Hemarthria altissima*). *Olant physiology*.69: 155-161.
- Vaughan. and Ord, B.G.(1991). Extraction of potential allelochemicals and their effects on root morphology and nutrient content. In.plant root growth an ecological perspective (Ed), D.Atkinson, *Black Well Scientific Publications London*, Po.399-421.
- Wilson, R.E and Rice, E.L. (1968). Allelopathy as expressed by *Helianthus annus* and its role in old-field succession. *Bulletin Torrey Botanical Club* 95:432-448

Identification of Allelochemicals from *Terminalia Chebula*

Table 1: Bioassay studies of aqueous leaf extract of *Terminalia chebula* on seed germination and seedling growth of *Cassia occidentalis*

Concentration %	Leaf Extract		
	Germination %	Plumule length (cm)	Radicle length (cm)
Control	70 ± 7.0	4.5 ± 0.4	3.5 ± 0.3
5	65 ± 6.2	4.2 ± 0.4	3.5 ± 0.3
10	60 ± 6.0	4.0 ± 0.3	3.2 ± 0.3
15	54 ± 5.0	3.8 ± 0.4	3.0 ± 0.2
20	50 ± 5.0	3.5 ± 0.3	2.7 ± 0.2

Table 2: Bioassay studies of aqueous bark extract of *Terminalia chebula* on seed germination and seedling growth of *Cassia occidentalis*

Concentration %	Bark Extract		
	Germination %	Plumule length (cm)	Radicle length (cm)
Control	76 ± 6.8	4.5 ± 0.4	3.5 ± 0.3
5	50 ± 5.0	3.0 ± 0.3	3.0 ± 0.2
10	45 ± 4.3	2.5 ± 0.3	2.7 ± 0.2
15	40 ± 3.8	2.0 ± 0.2	2.5 ± 0.2
20	36 ± 3.5	1.8 ± 0.2	2.0 ± 0.2

Table 3: Bioassay studies of aqueous leaf extract of *Terminalia chebula* on seed germination and seedling growth of *Crotalaria retusa*

Concentration %	Leaf Extract		
	Germination %	Plumule length (cm)	Radicle length (cm)
Control	60 ± 6.0	4.0 ± 0.4	3.5 ± 0.3
5	50 ± 5.0	3.0 ± 0.3	3.0 ± 0.3
10	45 ± 4.2	2.3 ± 0.2	2.7 ± 0.3
15	40 ± 4.0	2.0 ± 0.2	2.0 ± 0.2
20	30 ± 3.0	1.5 ± 0.1	1.4 ± 0.1

Table 4: Bioassay studies of aqueous bark extract of *Terminalia chebula* on seed germination and seedling growth of *Crotalaria retusa*

Concentration %	Bark Extract		
	Germination %	Plumule length (cm)	Radicle length (cm)
Control	60 ± 6.0	4.0 ± 0.4	3.5 ± 0.3
5	60 ± 5.7	3.5 ± 0.3	3.2 ± 0.3
10	55 ± 5.0	3.2 ± 0.3	3.0 ± 0.2
15	50 ± 5.1	3.0 ± 0.3	2.8 ± 0.3
20	46 ± 4.4	2.5 ± 0.2	2.5 ± 0.2

Table 5: Identification and content of phenolic acids in *Terminalia chebula* leaf and bark by gas chromatography

S.No	Phenolic acid	Relative retention time (min)	Content (µg/g Dr.wt)	
			Leaf	Bark
1	Hydro quinone	18.32	27.7	15.8
2	Salicylic acid	22.15	—	40.3
3	trans-cinnamic acid	22.84	20.3	—
4	Gentisic acid	26.64	153.8	3.2
5	Vanillic acid	29.79	34.4	4.2
6	Protocatechuic acid	31.56	—	5.7
7	Syringic acid	3.52	11.7	49.2
8	P-coumaric acid	34.46	—	30.1
9	trans-ferulic acid	38.35	67.2	4.2
10	Caffeic acid	39.54	—	2.1